

**SIGNALLING COMPOUNDS FOR USE IN METHODS OF
DETECTING HYDROGEN PEROXIDE**

CROSS REFERENCE TO RELATED APPLICATION

5 This application is a continuation-in-part of
applicant's co-pending U.S. Application Serial No.
10/371,053 filed on February 20, 2003.

10 FIELD OF THE INVENTION

 The present invention relates to methods and compounds
for detecting hydrogen peroxide. In particular, the present
invention relates to signalling compounds which react with
hydrogen peroxide to produce a detectable species. The
15 invention further relates to assay methods for detecting
hydrogen peroxide and for detecting peroxide-generating
systems. The invention further relates to boronic acid and
ester compounds, particularly fluorescent signalling
compounds, chemiluminescent signalling compounds and
20 dioxetane signalling compounds.

BACKGROUND OF THE INVENTION

1. Detection of Peroxide. Various methods are known for
detecting hydrogen peroxide and related peroxides.

25 a. Colorimetric Detection of Peroxide. Compounds which
react with hydrogen peroxide and a peroxidase to produce a
colored product include ABTS, 4-aminoantipyrine (Anal.
Letters, 26, 87 (1993)), and leuco dyes such as
Leucocrystal Violet. Numerous methods exist for detecting
30 hydrogen peroxide using a chromogen and a transition metal
compound.

b. Fluorescent Detection of Peroxide. Compounds which

react with hydrogen peroxide and a peroxidase to produce a fluorescent product include 2',7'-dichlorofluorescein, dihydrorhodamine 123 (Arch. Biochem. Biophys., 302(2), 348-55 (1993)) and N-acetylresorufin, (Chemical & 5 Pharmaceutical Bulletin, 49(3), 294-29, (2001)).

c. Chemiluminescent Detection of Peroxide. Acridinium esters and sulfonamides undergo a rapid oxidation reaction with hydrogen peroxide at alkaline pH to produce a flash of chemiluminescence (e.g. U.S. Patents 4,745,181, 4,946,958, 10 5,281,712 and 5,468,646). Lucigenin (9,9'-biacridinium dinitrate) is oxidized by hydrogen peroxide to produce chemiluminescence (Maskiewicz, et al., J.Am.Chem.Soc., 101, 5347-5354 (1979)).

Esters and amides of oxalic acid react with hydrogen 15 peroxide in the presence of a fluorescer to produce chemiluminescence. This reaction formed the basis of the well-known "light stick" technology used in novelty items.

Cyclic acylhydrazides including the amine-substituted compounds luminol and isoluminol, hydroxy-substituted 20 compounds and heterocyclic analogs react with hydrogen peroxide and a metal catalyst to produce chemiluminescence. Metal catalysts include heme, hexacyanoferrate and other transition metal ions including Cu(II) and Co(II).

U.S. Patent 5,545,834 describes the chemiluminescent 25 reaction of spiro-acridan compounds with hydrogen peroxide. The reaction is enhanced by the addition of horseradish peroxidase.

d. Enzymatic Detection of Peroxide. Various reagents have been developed for detection of peroxidase activity by 30 reaction of a peroxidase enzyme, a source of hydrogen

peroxide and an indicating reagent. These reagents therefore also serve to detect hydrogen peroxide. Color, fluorescence or chemiluminescence can be produced with use of the appropriate reagent. Chemiluminescent substrates
5 include amino- substituted cyclic acylhydrazides such as the well-known luminol and isoluminol (Anal. Chim. Acta, 170, 101-107, (1985)), heterocyclic acylhydrazides (M. Ii, et al., Biochem. Biophys. Res. Comm., 193(2), 540-5 (1993); U.S. Patent No. 5,324,835 and Y. Tominaga, et al.,
10 Tetrahedron Lett., 36, 8641-4 (1995)), and hydroxy-substituted phthalhydrazides (U.S. Patent No. 5,552,298).

Applicant's U.S. Patent Nos. 5,491,072, 5,523,212 and 5,593,845 disclose chemiluminescent N-alkylacridan-carboxylic acid derivatives which produce light upon
15 reaction with a peroxide and a peroxidase. Applicant's U.S. Patent No. 5,922,558 discloses a class of compounds containing an electron-rich double bond as chemiluminescent peroxidase substrates.

European Patent Specification EP0682254B1 discloses
20 assay methods in which a conjugate of an enzyme that generates hydrogen peroxide is used and the peroxide is detected by acridinium ester chemiluminescence.

Fluorescent substrates for peroxidase include 3-(4-hydroxyphenyl)propionic acid as disclosed in U.S.
25 Patent 6,040,150, 2-(4-hydroxyphenyl)acetic acid disclosed in Zaitzu and Ohkura, Anal. Biochem., 109, 109-113 (1980), homovanillic acid and tyramine (Y. Li, et al., Anal. Chim. Acta, (340), 159-168, (1997)), o-phenylenediamine and N,N'-dicyanomethyl-o-phenylenediamine (Li, et al., Microchem.
30 J., 53(4) , 428-436 (1996)), amide and carbamate

derivatives of p-aminophenol (M. Kawaguchi, et al.,
Bioluminescence and Chemiluminescence Perspectives for the
21st Century, A. Roda et al., Eds., Wiley & Sons,
Chichester, pp 508-511, (1999)), 3,4-dihydro-2(1H)-
5 quinoxalone and related derivatives (Li, et al., *Anal.*
Chim. Acta, 340(1-3), 159-168 (1997)), reduced forms of
fluorescein, rhodamine and other xanthine dyes and
fluorinated derivatives of the latter (U.S. Patent
6,162,931).

10 Chromogenic or color-forming substrates for peroxidase
include tetramethylbenzidine, chlorophenol red and 2,2'-
Azino-bis-(3-ethylbenzthiazoline-6-sulfonic acid (Ngo, T.
T. In *Immunochemistry of Solid-Phase Immunoassay*, Butler,
J. E. Ed., CRC: Boca Raton, 1991, pp 85-102.)).

15 2. Fluorescent and Colored Boronic Acid Sensors.
Fluorescent and colored compounds containing boronic acid
substituents for use in detection methods are disclosed in
U.S. Patents 4,496,722, and 4,659,817. The boronic acid
group complexing compounds coordinate to and bind pairs of
20 hydroxy, amine or thiol groups to form a fluorescent or
colored complex. The binding partner can be either a
carrier compound such as a buffer salt or it can be a
biological substance which is to be tagged. Representative
of the latter are cellular components. These methods differ
25 fundamentally from the methods of the present invention by
virtue of the unbound boronic acid compound being already
colored or fluorescent to the same extent as the bound
complex with the carrier or cellular component. Color or
fluorescence is not created during the conduct of the
30 methods.

The '772 and '817 patents also disclose peroxide assays in which the boronic acid substituted fluorescent or colored compounds are reacted with hydrogen peroxide to cleave the fluorescent or colored reporter moiety from the boronic acid group. Detection requires a separation step in order to measure the liberated reporter. Organic solvent extraction, release from a solid phase and filtration are disclosed as means to separate the released reporter from the reporter-boronic acid compound. Again, in contrast to the methods of the present invention, color or fluorescence is not created during the conduct of the methods.

Colored or fluorescent boronic acid complexing agents for use in detection of glycated blood proteins such as hemoglobin are disclosed in U.S. Patents 5,242,842, 5,506,144 and 5,739,318. The boronic acid groups coordinate to and bind pairs of hydroxy groups to form a colored or fluorescent complex. The complex is separated from unbound complexing agent and measured. Detection does not involve any peroxide. This mode of measurement differs from the present invention in requiring a separation and that the unbound boronic acid compound is already colored or fluorescent to the same extent as the bound protein complex.

PCT Publication WO 02/46752 discloses assays for polyhydroxyl compounds, e.g. glucose, using boronic acid-quencher conjugates which bind to and quench the fluorescence of appropriately substituted fluorescers. Binding of the fluorescer via the boronic acid group of the quencher is reversed in the presence of the polyhydroxyl compound which competes for binding with the boronic acid

group.

3. Phenylboronic acid Peroxidase Enhancers. U. S.

Patents 5,512,451 and 5,629,168 disclose phenylboronic acid compounds as enhancers of the peroxidase-catalyzed
5 chemiluminescent oxidation of luminol with hydrogen peroxide. In the methods disclosed therein the boronic acid compound promotes the reaction of the peroxidase in oxidizing luminol. Chemiluminescence is produced from an oxidized form of luminol and not from the boronic acid.
10 Arylboronic acid derivatives are disclosed as peroxidase enhancers in the chemiluminescent oxidations of acridan compounds in U.S. Patents 5,723,295 and 5,922,558.

None of the foregoing methods disclose the use of boronic acid or boronate ester signalling compounds in a
15 method for detecting and, when desired, quantifying the amount of hydrogen peroxide wherein the peroxide causes the formation of a detectable signal from the precursor signalling compound which does not itself possess the property being detected.

20

SUMMARY OF THE INVENTION

It is an object of the present invention to provide boronic acid or boronate ester signalling compounds which react with a source of hydrogen peroxide to produce a
25 detectable product having a detectable property, which compounds do not possess the property being detected.

It is an object of the present invention to provide compounds which react with a source of hydrogen peroxide to produce a colored, fluorescent, chemiluminescent or
30 bioluminescent product.

It is a further object of the present invention to provide signalling compounds used in methods for detecting a source of hydrogen peroxide.

It is a further object of the present invention to
5 provide signalling compounds for quantifying the amount of hydrogen peroxide by use of the above signalling compounds.

It is a further object of the present invention to provide signalling compounds for detecting enzymes which generate hydrogen peroxide by reaction of the enzyme with a
10 substrate for the enzyme and detecting the hydrogen peroxide generated by use of the above compounds.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 is a graph relating the amount of hydrogen
15 peroxide to the chemiluminescence intensity at 15 min emitted by 100 μ L of a reagent described in Example 10. The term S-B refers to the chemiluminescence signal (S) in Relative Light Units (RLU) in the presence of hydrogen peroxide corrected for background chemiluminescence (B) in
20 the absence of hydrogen peroxide.

Figure 2 is a graph relating the amount of glucose oxidase to fluorescence intensity according to Example 16. Samples containing various amounts of glucose oxidase were incubated with 0.1 M glucose for 30 min. The H_2O_2 produced
25 was assayed with a reagent containing the signalling compound 2-naphthylboronic acid to produce the fluorescent compound 2-naphthol.

Figure 3 is a graph relating the amount of glucose to fluorescence as described in Example 17. Samples containing
30 various amounts of glucose were incubated at room

temperature with glucose oxidase and the signalling compound 2-naphthylboronic acid. Fluorescence was measured after 30 min.

Figure 4 is a graph depicting the fluorescence spectra of a bis(benzothiazolyl)boronate ester signalling compound before (A) and after (B) reaction with urea peroxide.

Figure 5 is a graph depicting the grow in of fluorescence at 440 nm from the reaction of 4-methylcoumarin-7-boronic acid pinacol ester in 0.3 M tris buffer, pH 9.3 with 3 mM urea peroxide.

DESCRIPTION OF THE PREFERRED EMBODIMENTS

Definitions:

Alkyl - A branched, straight chain or cyclic hydrocarbon group containing from 1-20 carbons which can be substituted with 1 or more substituents other than H. Lower alkyl as used herein refers to those alkyl groups containing up to 8 carbons.

Alkenyl - A branched, straight chain or cyclic hydrocarbon group containing at least one C-C double bond and containing from 2-20 carbons. Lower alkenyl as used herein refers to those alkenyl groups containing up to 8 carbons.

Alkynyl - A branched or straight chain hydrocarbon group containing at least one C-C triple bond and containing from 2-20 carbons. Lower alkynyl as used herein refers to those alkynyl groups containing up to 8 carbons.

Analyte - A substance the presence or amount of which is to be measured in a sample by an assay. Analytes include organic and biological molecules to which a specific

binding partner having a specific binding affinity exists. Exemplary analytes include, without limitation, single stranded or double stranded DNA, RNA, DNA-RNA complexes, oligonucleotides, antibodies, antibody fragments, antibody-DNA chimeras, antigens, haptens, proteins, lectins, avidin, streptavidin and biotin. Other exemplary analytes also include hydrolytic enzymes, inhibitors of hydrolytic enzymes and dihydroxyaromatic compounds.

Aryl - An aromatic ring-containing group containing 1 to 5 carbocyclic aromatic rings, which can be substituted with 1 or more substituents other than H.

Biomedical analysis - Analyses of samples of biological origin for analytes of interest. The analyses can be immunoassays, western blots, northern blots, Southern blots, DNA hybridization assays, DNA sequence analysis, colony hybridizations, gene expression analysis, high throughput drug screening, detection of infectious agents or pathogens and the like.

Detectable signal - physical property resulting from measurement of the reaction product of the present reactions. The signal can be light produced by a chemiluminescent or bioluminescent reaction or by fluorescence. The signal can also be formation of color or a change of color or the change of absorption of ultraviolet or infrared radiation. The signal can also be a measurement of the molecular mass of the product or a specific signal in an NMR spectrum.

Halogen - Fluorine, chlorine, bromine or iodine atoms.

Heteroaryl or heteroaromatic - An aromatic ring-containing group containing 1 to 5 carbocyclic aromatic

rings in which at least one of the ring carbon atoms is replaced with a nitrogen, oxygen or sulfur atom and which can be substituted with 1 or more non-H substituents.

Luminescent - capable of emitting light when excited to
5 an electronic excited state. The light can be emitted either as fluorescence when decaying from a singlet excited state or as phosphorescence when decaying from a triplet excited state.

Peroxide - A compound containing an O-O bond,
10 preferably hydrogen peroxide or a complex of hydrogen peroxide such as urea peroxide, perborate or percarbonate.

Sample - A fluid containing or suspected of containing one or more analytes to be assayed. Typical samples which are analyzed by the chemiluminescent reaction method are
15 biological samples including body fluids such as blood, plasma, serum, urine, semen, saliva, cell lysates, tissue extracts and the like. Other types of samples include solvents, seawater, industrial water samples, food samples and environmental samples such as soil or water.


20 Source of hydrogen peroxide - a compound which is hydrogen peroxide or its salts such as sodium or potassium peroxide or hydrogen peroxide in complexed form such as urea peroxide, perborate salts and percarbonate salts.

Specific binding pair - Two substances which exhibit a
25 mutual binding affinity. Examples include antigen-antibody, hapten-antibody or antibody-antibody pairs, complementary oligonucleotides or polynucleotides, avidin-biotin, streptavidin-biotin, hormone-receptor, lectin-carbohydrate, IgG-protein A, nucleic acid-nucleic acid binding protein
30 and nucleic acid-anti-nucleic acid antibody.


Substituted - Refers to the replacement of at least one hydrogen atom on a group by a non-hydrogen group. It should be noted that in references to substituted groups it is intended that multiple points of substitution can be present unless clearly indicated otherwise.

Triflate - trifluoromethanesulfonate ester CF_3SO_3^- .

The present invention concerns methods using signalling compounds which react with a source of hydrogen peroxide to produce a detectable compound capable of producing a detectable signal. The signalling compounds comprise one or more boronic acid or boronate ester groups situated on a signalling moiety designated Sig.



In the signalling compounds each R is independently selected from hydrogen, alkyl and aryl groups and can be joined together as a straight or branched alkylene chain forming a ring or as an aromatic ring. The group Sig is not readily detectable until the boron-containing group is replaced by a hydroxyl group through the reaction of hydrogen peroxide. Thus, reaction of the signalling compound with hydrogen peroxide replaces the boron-containing substituent with a hydroxyl group creating the detectable moiety. The product Sig - OH is readily distinguished from the starting signalling molecule by the creation of a detectable property. The starting signalling molecule itself does not possess the detectable property or does so only to a very weak degree so that the product is easily distinguished and quantifiable.



Signalling compound Detectable product

5 Although the detectable product is depicted in the
 protonated form having a hydroxy group -OH, the detectable
 product can and in many cases will exist in the
 deprotonated form as the oxyanion or as an equilibrium
 mixture of protonated and deprotonated form. The exact form
 10 will be governed by factors such as reaction pH and
 electronic excitation which is known to ionize groups such
 as phenols by increasing their acidity in an electronic
 excited state. Both forms are considered to be within the
 scope of detectable product species in all embodiments of
 15 the invention.

Furthermore the present invention concerns methods of
 detecting hydrogen peroxide comprising reacting the
 signalling compounds with a source of hydrogen peroxide to
 produce a detectable product compound having a detectable
 20 property, detecting the product by measuring the detectable
 property and relating the detectable product to the
 hydrogen peroxide. The methods of the invention can be used
 in a quantitative fashion to determine the amount of
 hydrogen peroxide in a sample known or suspected to contain
 25 hydrogen peroxide. Such a method would comprise the steps
 of reacting the sample known or suspected to contain
 hydrogen peroxide with a signalling compound to produce a
 detectable product compound, detecting the detectable
 product compound by measuring a detectable property, and
 30 relating the detectable product to the amount of hydrogen

peroxide.

Detectable properties which can be used to detect and thereby differentiate the product from the signalling compound reactant broadly comprise any physical property which is altered by the reaction with peroxide and include color or absorption of light of any suitable wavelength from ultraviolet to visible to infrared, fluorescence, chemiluminescence, bioluminescence, molecular mass, and other means directly influenced by molecular structure including nuclear magnetic resonance frequency (NMR), especially ^1H and ^{13}C NMR. Preferred properties permit quantification of the amount of hydrogen peroxide by quantifying the amount or fraction of the detectable product which is formed and are selected from color, absorbance, fluorescence, chemiluminescence, and bioluminescence. The detectable property of the product compound preferably differs from that of the signalling compound by at least a factor of ten, more preferably by at least a factor of 100 and still more preferably by at least a factor of 1000. Preferred detectable properties are selected from fluorescence, chemiluminescence and bioluminescence. Importantly, detection of the product can be performed in the presence of the reactant signalling compound and does not require a separation.


Sources of hydrogen peroxide include hydrogen peroxide and its salts such as sodium or potassium peroxide, and peroxide in complexed form such as urea peroxide, perborate salts, percarbonate salts and percarboxylic acids and their salts. Biological sources of hydrogen peroxide are included as hydrogen peroxide is known to be produced in vivo in

leukocytes and certain antibodies. Sources of hydrogen peroxide also encompasses enzymatic generation systems as discussed below.

5 Signalling compounds useful in the methods of the invention comprise one or more boronic acid or boronate ester groups situated on a signalling moiety designated Sig. The group Sig can be any organic group capable of being detected as described above. Because of the near universality of detection of organic compounds by
10 techniques such as NMR spectroscopy and mass spectrometry the nature of the Sig group encompasses essentially any non-polymeric organic group. Aliphatic, unsaturated, aromatic and heterocyclic groups can be used as the Sig group. The detectable property when using mass
15 spectrometric detection is the molecular mass of the product which of course differs from that of the starting signalling compound by at least 27 mass units up to about 200 mass units depending on the nature of the groups R^5 and R^6 . The group Sig is limited only by the mass resolution of
20 the measurement and the ability to detect the molecular ion. Sig groups having a molecular mass up to about 2000 will be detectable.

 Detection by NMR spectroscopy allows detection of any reaction product Sig-OH which can be distinguished from the
25 corresponding signalling compound. The detection can be performed by ^{13}C NMR analysis of the resonance frequency of the carbon undergoing the C-B to C-O bond change. Monitoring the change is most conveniently performed in ^1H -decoupled mode so that the signal will be a singlet peak in
30 most cases. The detection can also be performed by ^1H NMR

analysis of the resonance frequency of a hydrogen atom on or near the carbon undergoing the C-B to C-O bond change. By near is meant that the hydrogen atom is substituted on another carbon atom or other atom within 1-3 atoms of the carbon undergoing the C-B to C-O bond change. Some examples of hydrogen atoms detectable by the present invention and useful in a qualitative or quantitative method for detecting the presence or amount of hydrogen peroxide are shown below.




Signalling compounds detectable by NMR techniques when reacted in accordance with the present invention are preferably non-polymeric organic compounds wherein the Sig group has a molecular mass less than about 2000. Desirably the group Sig will not contain other atoms or groups which are chemically similar to the particular nucleus being detected and thereby obscure its detection. Quantification is readily achieved using NMR-based detection. A sample containing hydrogen peroxide is reacted with a signalling compound in an amount at least equal to the amount of the peroxide. The entire sample or a known fraction thereof is analyzed and the extent of conversion to the reaction

product Sig-OH determined from the magnitude of the selected characteristic resonance signal.

A preferred class of groups for Sig comprises substituted and unsubstituted aromatic or heteroaromatic ring groups. Signaling compounds in this class generally possess a chromophore or fluorophore when converted to the product by reaction with a source of hydrogen peroxide and can therefore be used with an absorption or fluorescence detection scheme.

One group of signalling compounds comprise those boronic acid or boronate ester compounds which, upon conversion to the phenolic product Sig - OH through reaction with a source of hydrogen peroxide, become capable of generating chemiluminescence. Among the signalling compounds of the present invention which are converted to chemiluminescent products are a novel family of 1,2-dioxetane compounds. The previously unknown boronic acid or boronate ester substituted dioxetanes are of formula I:




wherein A¹ - A³ represent organic groups having from 1-20 carbon atoms and Ar is an aromatic or heteroaromatic ring group, wherein A¹ and A² or A¹ and A³ or A³ and Ar can be combined to form a ring, and each R is independently selected from hydrogen, alkyl and aryl groups and can be joined together as a straight or branched alkylene chain forming a ring or as an aromatic ring. All such groups A¹-A³, Ar and R can be substituted with non-hydrogen atoms.

These compounds have been unexpectedly discovered to react

with hydrogen peroxide to produce chemiluminescence.

A group of preferred 1,2-dioxetane compounds for detection of hydrogen peroxide by chemiluminescence have formula II:

5




II

wherein R¹ is is an organic group having from 1-20 carbon atoms which can be substituted with non-hydrogen atoms and can be combined with R² or R³, R² is an aromatic or heteroaromatic ring group which can include additional substituents selected from halogens, alkyl, substituted alkyl, alkoxy, substituted alkoxy, carbonyl, carboxyl, amino and alkylamino groups, R³ and R⁴ are independently selected from acyclic and cyclic organic groups containing from 3-20 carbon atoms and which can be substituted with heteroatoms and R⁵ and R⁶ are independently selected from hydrogen, alkyl and aryl groups and and can be joined together as a straight or branched alkylene chain forming a ring or as an aromatic ring. In another group of compounds of formula II, R³ and R⁴ are combined together in a cyclic or polycyclic alkyl or alkenyl group which is spiro-fused to the dioxetane ring and contains 6 to 20 carbon atoms and which can include additional non-hydrogen substituents. A preferred polycyclic alkyl group is a substituted or unsubstituted adamantyl group.

A preferred group of compounds are boronic acids wherein R⁵ and R⁶ are both hydrogen atoms. Another preferred group of compounds contain an R² group selected from substituted or unsubstituted phenyl and substituted or

unsubstituted naphthyl groups. In another preferred group of compounds R^1 is a lower alkyl group which can be substituted with non-hydrogen atoms. In another preferred group of compounds R^3 and R^4 are each branched alkyl or cycloalkyl groups having from 3-20 carbon atoms. In another preferred group of compounds R^3 and R^4 are combined together as a spiro-fused polycyclic alkyl or alkenyl group and is more preferably an adamantyl group or a substituted adamantyl group having one or more substituent groups selected from halogens, alkyl, substituted alkyl, alkoxy, substituted alkoxy, carbonyl, carboxyl, phenyl, substituted phenyl, amino and alkylamino groups covalently bonded thereto.



Dioxetane compounds of formula I and II are converted by reaction with a source of hydrogen peroxide to the corresponding hydroxy-substituted dioxetane which produces chemiluminescence rapidly when in the deprotonated state.





A large number of species are known to function in the method of the present invention and are encompassed within the class of compounds embodied by formula I. This knowledge is predicated in part on the knowledge that once a hydroxy-substituted dioxetane is formed by reaction of a boronic acid or boronate ester dioxetane derivative with peroxide, it will spontaneously decompose when deprotonated to produce light. Many such triggerable dioxetanes are known. Particular embodiments of compounds of formula I and II which react according to the present invention to

produce chemiluminescence comprise those boronic acid or boronate ester dioxetanes which are converted by hydrogen peroxide to known hydroxy-substituted dioxetanes. These include, without limitation

5





10 wherein R⁷ is a cyclic or polycyclic alkyl or alkenyl group,



15



wherein Y is a substituent group selected from halogens, alkyl, substituted alkyl, alkoxy, substituted alkoxy, carbonyl, carboxyl, phenyl, substituted phenyl, amino and alkylamino groups,

20







wherein Z is a substituent group selected from halogens, alkyl, substituted alkyl, alkoxy, substituted alkoxy, carbonyl, carboxyl, phenyl, substituted phenyl, amino and alkylamino groups,









25





30



wherein n is from 1 to 3







5



wherein $n = 1$ or 2 , R is selected from alkyl and benzyl and R' is selected from alkyl, phenyl and substituted phenyl; T is selected from alkyl, cycloalkyl and polycycloalkyl groups any of which can be substituted with non-hydrogen atoms, Ak is a branched alkyl or cycloalkyl group which can be substituted with non-hydrogen atoms,

15




wherein W is O or S ;

20




particularly where the group R^1 is lower alkyl or halogenated alkyl such as CH_3 , CD_3 , or CH_2CF_3 and Y is 4-chloro or H

25



and




30


wherein W is O or S ;

and boronate ester derivatives thereof as described above.

In addition to the foregoing structures, boronate ester analogs of these compounds are also useful in the practice of the present methods. In place of the $B(OH)_2$ group, the dioxetane compounds can contain a group having the formula $B(OR)_2$ wherein R can be an alkyl, specially a lower alkyl, e.g. methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, t-butyl and the like or an aryl ring group such as phenyl. The R groups can be joined together in a ring e.g.




Another group of signalling compounds of formula I comprise dioxetane compounds substituted with a sulfur atom on the dioxetane ring. These include, without limitation



III

and



wherein R^1 , R^2 , R^3 , R^4 , R^5 , and R^6 are as described above, and R^7 is a cyclic or polycyclic alkyl or alkenyl group any of which can be substituted with non-hydrogen atoms.


Particularly, signalling compounds of formula III comprise compounds in which R^1 is an organic group having from 1-20

carbon atoms which can be combined with R^2 or R^3 , R^2 is an aromatic or heteroaromatic ring group which can include additional substituents selected from halogens, alkyl, substituted alkyl, alkoxy, substituted alkoxy, carbonyl, carboxyl, amino and alkylamino groups, R^3 and R^4 are independently selected from acyclic and cyclic organic groups containing from 3-20 carbon atoms and which can be substituted with heteroatoms, and R^5 and R^6 are independently selected from hydrogen, alkyl and aryl groups and can be joined together as a straight or branched alkylene chain forming a ring or as an aromatic ring.

Preferred embodiments include without limitation



wherein Y is a substituent group selected from hydrogen, halogens, alkyl, substituted alkyl, alkoxy, substituted alkoxy, carbonyl, carboxyl, phenyl, substituted phenyl, amino and alkylamino groups,




25 wherein Z is a substituent group selected from hydrogen, halogens, alkyl, substituted alkyl, alkoxy, substituted alkoxy, carbonyl, carboxyl, phenyl, substituted phenyl, amino and alkylamino groups.

Another aspect of the present invention is the use of a signalling compound including the compounds described above

in a method to produce chemiluminescence by reaction with hydrogen peroxide. Reaction of the compound with hydrogen peroxide in an aqueous solution produces easily detected chemiluminescence when the reaction is conducted at

5 alkaline pH. The reaction can be conducted optionally in the presence of a chemiluminescence enhancer.

In a preferred method of producing chemiluminescence, compound I or II is reacted with a peroxide in an alkaline solution with a pH of at least about 8 to produce
10 chemiluminescence which commences upon reaction of the peroxide and compound I or II according to the reaction scheme.



Reaction of the boronic acid or boronate ester-substituted dioxetane with hydrogen peroxide is believed to form a hydroxyaryl or aryloxy substituted dioxetane
20 intermediate by replacing the carbon-boron bond with a carbon-oxygen bond. Under conditions in which of the phenolic OH group is at least partly ionized, decomposition of the dioxetane ensues with emission of light. The base or alkaline pH-triggered decomposition of hydroxyaryl
25 substituted dioxetanes is well known in the art of chemiluminescence. Examples of such triggerable dioxetanes with various structural modifications are described in for example, U.S. Patents 5,886,238, 6,036,892, 6,284,899, 6,410,751, 5,650,525, 5,731,445, 5,877,333, 5,929,254,
30 6,218,135, 6,228,653, 5,013,827, 5,068,339, 5,652,345,

5,770,743, 5,132,204, 5,248,618, 5,603,868, 5,712,106,
6,107,036, 4,952,707, 5,089,630, 5,112,960, 5,220,005,
5,326,882, 5,330,900, 5,538,847, 5,543,295, 5,582,980,
5,591,591, 5,625,077, 5,679,802, 5,707,559, 5,773,628,
5 5,783,387, 5,831,102, 5,840,919, 5,843,681, 5,851,771,
5,869,699, 5,869,705, 5,871,938, 5,981,768, 6,022,964,
6,063,574, 6,132,956, 6,133,459, 6,140,495, 6,355,441 and
6,461,876. The boronic acid and boronate ester analogs of
all of these dioxetanes are considered to fall within the
10 scope of the present invention and are expressly included
as part of the disclosure of the invention.


Chemiluminescence can also be produced from the above
signalling compounds by reaction with a source of hydrogen
peroxide followed by reaction of the hydroxy-substituted
15 dioxetane in an organic solvent with a strong base such as
hydroxide ion or fluoride ion. Polar aprotic solvents such
as DMSO, DMF and acetonitrile are preferred for producing
highest intensity light.

Use of the above triggerable dioxetane signalling
20 compounds of the present invention results in formation of
a detectable product that can also be detected on the basis
of the change in molecular mass and by ^1H or ^{13}C NMR as
described above. Moreover it is recognized that the
chemiluminescent reaction of the triggerable dioxetane
25 signalling compounds results in a further fragmentation of
the product in the process of producing chemiluminescence.
It is recognized that the fragmentation or cleavage
products can also be detected and/or quantified by
molecular mass and by ^1H or ^{13}C NMR as described above.
30 Other reaction products described below that are detectable


by a chemiluminescent reaction also form subsequent products as the result of a fragmentation reaction or an oxidative dimerization process. These further products have unique molecular mass and NMR properties which can serve as the basis of detection.

Substances that are useful as chemiluminescence enhancers are those already known in the art of chemiluminescence. Included among these are quaternary onium salts alone or in combination with anionic surfactants such as alkyl sulfate salts and alkyl sulfonate salts. Quaternary onium salts include quaternary ammonium and phosphonium salts such as cetyltrimethylammonium bromide, dicationic surfactants as described in U.S. Patent 5,451,437, polymeric phosphonium salts as described in U.S. Patent 5,393,469, polymeric ammonium salts as described in U.S. Patent 5,112,960, polymeric mixed phosphonium and ammonium salts and mixtures of polymeric ammonium salts and additives as described in U.S. Patent 5,547,836.


A representative procedure for the synthesis of a boronic acid-substituted dioxetane useful as a chemiluminescent signalling compound is depicted below.



Additional chemiluminescent signalling compounds of the invention have the formulas below




wherein Z is selected from O, S and NR^8 , wherein R^8 is H or $\text{Si}(\text{R}^9)_3$, R^9 is C_1 - C_6 alkyl or phenyl, and X represents one or two halogen substituents, preferably iodine, bromine or chlorine atoms. Reaction of the above signalling compounds with hydrogen peroxide replaces the boronic acid (or ester) substituent with a hydroxy group. Hydroxy-substituted indole compounds are described in e.g. U.S. Patent 5,589,238 as undergoing a spontaneous chemiluminescent reaction with ambient oxygen.




The above indolyl, benzofuranyl and benzothiophenyl boronic acid derivatives can also be used in embodiments where color is measured since the indigo-like dimerized products are highly colored. Moreover, the aforementioned U.S. Patent 5,589,238 as well as European Patent Specification EP 0476930 B1 describe that this reaction also produces additional hydrogen peroxide, presumably through reduction of oxygen. Use of an indolyl, benzofuranyl or benzothiophenyl boronic acid signalling compound of the present invention will allow detection with

greater sensitivity since the hydrogen peroxide will be recycled in the reaction. Stated another way, each molecule of hydrogen peroxide is capable of converting many molecules of signalling compound to detectable product.

5 Other chemiluminescent signalling compounds of the invention have the formula below




wherein LG is a leaving group and R^{10} and R^{11} are hydrogen, C_1 - C_4 alkyl or are combined as an alkylene ring. Exemplary signalling compounds include the carboxylic acid, ester and thioester derivatives.



In the structure above, R is a substituted or unsubstituted alkyl or aryl group, and AMP designates adenosine monophosphate. Reaction of the boronic acid or ester moiety with

25 hydrogen peroxide will produce the molecule known as firefly luciferin. This compound is known to react with the firefly luciferase enzyme to produce bioluminescence. It is also known to produce chemiluminescence upon treatment with strong base by an oxidative process involving molecular

30 oxygen.



Embodiments in which the leaving group is the AMP ester and the luciferase enzyme is used constitute a use of the present invention in a bioluminescent method to detect a source of hydrogen peroxide. Such a method comprises

- 15 a) reacting a signalling compound with a source of hydrogen peroxide to produce a detectable product capable of undergoing a bioluminescent reaction;
- b) reacting the detectable product with an enzyme and any necessary co-factors for the enzyme to produce
- 20 bioluminescence; and
- c) detecting the bioluminescence.

The reaction can be conducted in different experimental modes. In one mode the carboxylic acid precursor can be reacted with peroxide in a first step to replace the boronic acid or ester with the hydroxyl group to produce


25 the luciferin. The luciferin thus formed is then subjected to the art-known conditions for eliciting bioluminescence, reaction with firefly luciferase, Mg^{+2} and ATP in a buffer. Since these two steps have differing pH optima, each step

30 can be conducted at its optimum pH or they can be conducted

at a single pH conducive to both reactions. In another mode of reaction the peroxide reaction is conducted concurrently with the luciferase/ATP reaction. reagents specifically identified for this purpose in U.S. Patents 5,618,682,
 5 5,650,289 and 5,814,471.

Another group of signalling compounds capable of being detected by chemiluminescence have the formula:

10




15

wherein R^5 and R^6 are as described above. Reaction of the compound above with a source of hydrogen peroxide converts the boronic acid or ester compound into the known hydroxyaryl hydrazide compound. The latter are known to undergo chemiluminescent reaction in aprotic solvent by reaction with strong base or in aqueous alkaline solution with additional peroxide and optionally with a transition metal catalyst or a peroxidase enzyme.

20


Another group of signalling compounds comprise those boronic acid or boronate ester signalling compounds which are converted to a fluorescent phenolic product Sig - OH through reaction with hydrogen peroxide. Among the signalling compounds of the present invention which are
 25 converted to fluorescent products are substituted and unsubstituted naphthaleneboronic acids and esters,

30




as well as the anthracene, phenanthrene and pyrene analogs,
compounds of the formula which produce a coumarin or
umbelliferone product,

5




benzothiazole boronic acids and esters of the formula

10



and

15




wherein Z is C-C double or triple bond or aromatic ring and
n is 1 or 2, R⁹ is a cyano, imine or carbonyl group, and
compounds which generate other fluorescers including

20

resorufin and fluorescein and having the structures below.

25




It should be recognized that the signalling compounds
described above which produce a chemiluminescent product
also can be detected by fluorescence. This situation arises

because the emitter in a chemiluminescent reaction is, in nearly all cases, also fluorescent.


The above-mentioned signalling compounds which are converted to a fluorescent phenolic product Sig - OH are useful in a method for detecting a source of hydrogen peroxide comprising:

- a) reacting a signalling compound with a source of hydrogen peroxide to produce a detectable product Sig - OH capable of detection by fluorescence;
- 10 b) irradiating the detectable product with light of a first wavelength; and
- c) detecting light emitted as fluorescence from the detectable product at a second wavelength different from the first wavelength.

15 Another group of signalling compounds comprise those boronic acid or boronate ester signalling compounds which are converted to a colored phenolic product Sig - OH through reaction with hydrogen peroxide. Among the signalling compounds of the present invention which are
20 converted to colored products include, without limitation, the exemplary structures depicted below.



and




5

Numerous other colored compounds containing a phenol moiety are known in the art of colorimetric assays. Signalling compounds based on these structures but containing a boronic acid or boronate ester moiety in place of the hydroxyl group are explicitly considered to be within the scope of the invention.

The above-mentioned signalling compounds which are converted to a colored phenolic product Sig - OH are useful in a method for detecting a source of hydrogen peroxide comprising:

- a) reacting a signalling compound with a source of hydrogen peroxide to produce a detectable product Sig - OH capable of detection by its color; and
- b) detecting the formation of the detectable product by the formation or change of color at a suitable wavelength or range of wavelengths.

Boronic acid and boronate ester compounds useful as signalling compounds as described herein are generally prepared by reaction of a boronating agent with an aryl halide or aryl triflate and a metal catalyst.



Y = Cl, Br, I, OSO₂CF₃

X = H, OR, B(OR)₂

30

The metal catalyst is preferably a palladium compound used in the presence of a phosphorus compound capable of acting as a ligand on palladium. Representative palladium catalysts useful for effecting carbon-boron bond formation are well known in the art of synthetic organic chemistry and include divalent compounds PdL_2 with labile ligands L selected from carboxylate esters, halogens and ketones and include palladium acetate, palladium chloride, palladium bis(dibenzylideneacetone) $\text{Pd}(\text{dba})_2$ and $\text{Pd}_2(\text{dba})_3$.

Phosphine ligands include PPh_3 , PMe_3 , PEt_3 , Pt-Bu_3 , $\text{Pt-Bu}_2\text{Me}$, $\text{Pt-Bu}_2\text{Et}$, $\text{P}(\text{cyclohexyl})_3$, BINAP and mixed alkylarylphosphines such as DPPE, DPPF, DPPB and DPPP. Bases include KHPO_4 , K_2CO_3 , Na_2CO_3 , CsCO_3 , CsF , Et_3N and alkoxide salts such as sodium t-butoxide. Solvents useful in this step include toluene, benzene, THF, DME, diglyme and alcohol solvents including t-butanol and t-amyl alcohol.

Commonly used boronating agents include trialkyl borates such as trimethylborate, triethylborate and triisopropyl-borate, diboron compounds including bis(pinacolato)diboron, bis(neopentylglycolato)diboron.

The aromatic component is generally a bromide, iodide or triflate derivative of an aromatic hydrocarbon such as benzene, naphthalene, anthracene or pyrene or an aromatic heterocycle such as pyridine, quinoline, acridine, pyrrole, furan, benzofuran, thiophene, benzothiophene, oxazole, benzoxazole, thiazole, benzothiazole, xanthene, thioxanthene and phenothiazine any of which can be further substituted or unsubstituted.

Other methods of synthesizing boronic acids are known.

Reaction of aryl bromide compounds with n-BuLi in THF at-
78 °C produces an aryllithium compound which is reacted in
situ with trimethyl borate or other trialkyl borates to
produce the arylboronate ester. Hydrolysis to the

- 5 arylboronic acid normally ensues during the reaction workup
when trimethyl borate is employed.

Likewise Grignard reagents can be utilized in place of
the aryllithium intermediates mentioned above in a method
of synthesizing arylboronic acids and their esters.

- 10 Reaction of the aryl Grignard reagent with a trialkyl
borate in ether solvent produces the arylboronate ester.

- Arylboronic acids and esters can also be prepared from
an aromatic ring compound in a process using mercury-
containing intermediates. Organomercury derivatives are
15 converted to arylboronic acids (Q. Zheng, et al.,
Heterocycles, 37, 1761-72. (1994); N. Garg, et al., J. Am.
Chem. Soc., 124, 13179 (2002)). For example, reaction of an
indole compound with $\text{Hg}(\text{OAc})_2$ produced the organomercurial
 $\text{Ar-Hg}(\text{OAc})$. Reaction of the latter with borane (BH_3 -THF)
20 followed by hydrolysis produced an indole-3-boronic acid.

- Arylboronate esters can also be prepared from arenes
according to the methods disclosed in Ishiyama, et al., J.
Am. Chem. Soc., 124, 390-1 (2002). Reaction of an arene
with an iridium catalyst with a bipyridyl or phenanthroline
25 ligand and a dialkylborane or diboron compound effects
direct borylation of the aromatic ring of the arene.

- Reactions of signalling compounds of the present
invention with peroxide are carried out in solution such as
an organic solvent or an aqueous buffer or mixtures
30 thereof. The reaction solution can be in contact with the

surface of a solid support such as a bead, tube, membrane or microwell plate. Reactions involving the use of enzymes are conveniently performed in a buffer. Suitable buffers include any of the commonly used buffers capable of
5 maintaining a pH in the range of about 6 to about 10 for example, phosphate, borate, carbonate, tris(hydroxymethyl-amino)methane ("tris"), glycine, tricine, 2-amino-2-methyl-1-propanol ("221"), diethanolamine and the like. The preferred method of practicing the invention in this regard
10 is determined by the intended use.

The amount of time used for performing the reaction of the signalling compound with the source of hydrogen peroxide can vary over a wide range. Under conditions of alkaline solution reaction can be rapid, proceeding to
15 completion in minutes. Under conditions of enzymatic generation of peroxide reaction times of hours can be used. Longer reaction times are not harmful even when the reaction is completed quickly. Conversely, short reaction times can be used even when it would not permit consumption
20 of all of the peroxide.

Embodiments involving fluorescent or chemiluminescent products require the detection of light emission. Light emitted by the present method can be detected by any suitable known means such as a luminometer, x-ray film,
25 high speed photographic film, a CCD camera, a scintillation counter, a chemical actinometer or visually. Each detection means has a different spectral sensitivity. The human eye is optimally sensitive to green light, CCD cameras display maximum sensitivity to red light, x-ray films with maximum
30 response to either UV to blue light or green light are

available. Choice of the detection device will be governed by the application and considerations of cost, convenience, and whether creation of a permanent record is required.

Signalling compounds of the present invention which
5 produce light by means of fluorescence, chemiluminescence or bioluminescence typically emit light over a 100-200 nm wide band of emission, which exhibits a maximum intensity at wavelengths in the near ultraviolet to the visible region of the electromagnetic spectrum. Typical wavelengths
10 of maximum intensity λ_{max} in the range of 350-750 nm. It is contemplated that signalling compounds bearing a covalently linked fluorophore could undergo intramolecular energy transfer resulting in emission at longer wavelengths from the excited state of the fluorophore.

15 In a further embodiment, luminescent energy acceptors can be employed to shift the maximum emission to longer wavelengths (red-shifting) and/or to increase the quantity of luminescence emitted. Various techniques for red-shifting emission are known in the art of chemiluminescent
20 reactions and assays. Covalently linked fluorophores are one example. Fluorescers can alternatively be added to the reaction solution as separate species. Fluorescers can be linked to a polymer or associated with a micelle or polymer in order to bring the fluorescer in close contact to the
25 compound. Suitable energy transfer agents have an excited state at an energy level which overlaps that of the excited reaction product to permit the transfer of excitation energy and a fluorescent excited state which may or may not be the same as the excited state which overlaps that of the
30 donor. Energy transfer agents useful for effecting singlet-

singlet energy transfer are well known in the art of fluorescence. Energy transfer agents useful for effecting triplet-singlet energy transfer are also known in the art and usually possess at least one metal atom or other heavy atoms such as bromine or iodine atoms. Typical examples are 9,10-dibromoanthracene (DBA), sulfonated derivatives of DBA and $\text{Ru}(\text{bpy})_3^{2+}$. Fluorescent energy transfer agents are evaluated empirically by comparing the intensity or wavelength of chemiluminescence produced in a reaction of a peroxide and a signalling compound as described above in the presence and absence of the agent.

Embodiments involving the formation of colored products require the detection of color or the absorption of light. Detection can be by any suitable known means. The simplest is visual observation of color development or color change. Photographic film or a digital camera can also be employed for this purpose. Embodiments requiring quantitative measurement of color will best be performed by spectrophotometry. Choice of the detection device will be governed by the application and considerations of cost, convenience, and whether creation of a permanent record is required.

The present invention includes the use of the methods and signalling compounds for detecting hydrogen peroxide in an assay procedure. Reaction of a signalling compound of the invention with a source of hydrogen peroxide produces the phenolic product which is detected by the appropriate means, i.e. color, absorbance, fluorescence or luminescence as described above. The magnitude of the signal is then related to the amount of peroxide present. Quantitative

relationships are readily established by constructing calibration curves for any given signalling compound and a set of peroxide standards of known concentration. Uses for detection of hydrogen peroxide include the monitoring of biological or cellular processes, detection of antibodies which generate hydrogen peroxide, and the analysis of peroxide contamination in peroxide-forming solvents. Several of the signalling compounds of the present invention when dissolved in ether solvents such as p-dioxane rapidly give rise to detectable product, indicating the presence of peroxide contaminant.

In a further embodiment the present invention also relates to the use of these methods for detecting an enzyme which produces peroxide such as an oxidase enzyme or a dehydrogenase enzyme. It is well known that various enzymes in the class of oxidase enzymes produce hydrogen peroxide as a byproduct of the reaction which oxidizes its substrate. Known oxidase enzymes include galactose oxidase, glucose oxidase, cholesterol oxidase, amine oxidase, various amino acid oxidases, polyphenol oxidase, xanthine oxidase, uricase, alcohol dehydrogenase, lactate dehydrogenase, malate dehydrogenase, glyceraldehyde-3-phosphate dehydrogenase, glycerol dehydrogenase, and glucose-6-phosphate dehydrogenase. In practice, the oxidase enzyme is reacted with a substrate for the oxidase enzyme to produce hydrogen peroxide. Either concurrently or after a suitable time period, the accumulated hydrogen peroxide is reacted with a signalling compound of the invention to produce the phenolic product which is detected by the appropriate means as described above. Further the oxidase

or dehydrogenase enzyme can be present as a conjugate to a biological molecule or a member of a specific binding pair in an assay for an analyte.

An important use of the present methods in biomedical
5 and other analysis is for detecting the presence or amount of an analyte in an assay procedure by a chemiluminescent, fluorescent, bioluminescent or color-forming reaction. One format comprises using an oxidase or dehydrogenase enzyme as a label on a specific binding pair member. An example is
10 an enzyme-linked immunoassay, such as the so-called enzyme-linked immunosorbent assay or ELISA. Such assays fall broadly into two categories. Competitive assays feature an immunological binding of a specific antibody with the analyte and an analyte analog, e.g. a detectably labeled
15 analyte molecule. Sandwich assays result by the sequential or simultaneous binding of two antibodies, one of which is detectably labeled, with the analyte. The detectably labeled binding pair so formed is assayed with the compounds and methods of the present invention. When the
20 detectable label is the oxidase enzyme, it is detected directly. When the detectable label is a member of another specific binding pair, e.g. a hapten, a conjugate of its binding partner with an oxidase is reacted first and the oxidase then detected in accordance with the present
25 methods. Measurement can be performed with enzyme-labeled species in a solution or attached to a solid surface or support including beads, tubes, microwells, magnetic particles, test strips, membranes and filters such as are in common use in the art. Other exemplary uses are the
30 detection of proteins by the technique of Western blotting

and nucleic acids by the use of enzyme-labeled nucleic acid probes including Southern blotting, northern blot analysis of RNA, and DNA sequencing.


Detection of the oxidase-labeled specific binding pair
5 formed in the assay is conducted by supplying a substrate for the oxidase enzyme and, either concurrently or subsequently, supplying a signalling compound of the invention. Reaction of the oxidase with its substrate produces hydrogen peroxide. The hydrogen peroxide reacts
10 with the signalling compound to produce the phenolic product which is detected by the appropriate means as described above.

The method comprises the steps of contacting a sample suspected of containing the analyte with a signalling
15 compound of the present invention and a source of hydrogen peroxide, and detecting the phenolic product by means of chemiluminescence, fluorescence, bioluminescence or color in a qualitative method. If quantitation is desired, the amount of phenolic product formed is related to the amount
20 of the analyte. The relationship between the detectable signal and amount of analyte can be easily discerned by constructing a calibration curve with known amounts of the analyte. The signalling compound is typically used in a concentration of about 10^{-8} M to about 10^{-2} M, preferably
25 between about 10^{-6} M and about 10^{-3} M. Typical samples which can be analyzed by the present methods are body fluids such as blood, plasma, serum, urine, semen, saliva, sputum, cerebrospinal fluid and the like.

The present invention also includes embodiments wherein
30 the signalling compound contains a labelling group other

than the boronic acid group which permits labeling on a substance to be detected. The labeling group comprises a group having the formula -L-RG wherein L is a linker and RG is a reactive group.

5




The group L is a linking group which can be a bond or another divalent or polyvalent group, the group RG is a reactive group which enables the signalling compound to be
10 linked to another compound.

The linking group can be a bond, an atom, or a straight, or branched chain of atoms some of which can be part of a ring structure. The substituent usually contains from 1 to about 50 non-hydrogen atoms, more usually from 1
15 to about 30 non-hydrogen atoms. Atoms comprising the chain are selected from C, O, N, S, P, Si, B, and Se atoms, preferably from C, O, N, P and S atoms. Halogen atoms can be present as substituents on the chain or ring. Typical functional groups comprising the linking substituent
20 include alkylene, arylene, alkenylene, ether, peroxide, carbonyl as a ketone, ester, carbonate ester, thioester, or amide group, amine, amidine, carbamate, urea, imine, imide, imidate, carbodiimide, hydrazine, diazo, phosphodiester, phosphotriester, phosphonate ester, thioether, disulfide,
25 sulfoxide, sulfone, sulfonate ester, sulfate ester, and thiourea groups.

The reactive group RG is an atom or group whose presence facilitates bonding to another molecule by covalent attachment or physical forces. In some
30 embodiments, attachment of a signalling compound of the

present invention to another compound will involve loss of one or more atoms from the reactive group, e.g. when the reactive group is a leaving group such as a halogen atom or a tosylate group and the signalling compound bonds to another compound by a nucleophilic displacement reaction. In other embodiments, attachment of a signalling compound to another compound by covalent bond formation will involve reorganization of bonds within the reactive group as occurs in an addition reaction such as a Michael addition or when the reactive group is an isocyanate or isothiocyanate group. In still other embodiments, attachment will not involve covalent bond formation, but rather physical forces in which case the reactive group remains unaltered. By physical forces is meant attractive forces such as hydrogen bonding, electrostatic or ionic attraction, hydrophobic attraction such as base stacking, and specific affinity interactions such as biotin-streptavidin, antigen-antibody and nucleotide-nucleotide interactions.



The structure above bearing a biotin moiety as a reactive group exemplifies a compound capable of specific binding interactions by non-covalent means. Numerous specific linking groups and reactive groups are listed in U.S. Patent 6,126,870.


The present invention also includes embodiments wherein the signalling compound is provided as a label on a

substance to be detected. By this is meant conjugates of a substance which is to be detected and at least one signalling compound IV bearing a labeling substituent.

In order to more fully describe various aspects of the present invention, the following examples are presented
5 which do not limit the scope of the invention in any way.


EXAMPLES

1. Synthesis of a Dioxetaneboronic Acid.



1

5 A mixture of 143.5 g of TiCl_3 and 17.64 g of LiAlH_4 was formed in 1 L of cooled anhydrous THF under Ar by slow addition of the LiAlH_4 to the reaction system. A black
10 mixture resulted. Triethylamine (130 mL) was added and the mixture heated to reflux. After 1 hour at reflux a solution of 20.62 g of methyl 3-bromobenzoate and 44.02 g of adamantanone in 200 mL of anhydrous THF was added through a dropping funnel over 30 min. The reaction mixture was cooled
15 and carefully poured into a solution of 6 L of water and 300 mL of triethylamine. The mixture was extracted with 7 x 1 L of ethyl acetate and discarded. The ethyl acetate extracts were combined, dried over Na_2SO_4 , filtered and evaporated to dryness leaving a white solid. The solid was
20 washed several times with hexane and discarded. The hexane solutions were combined, concentrated in volume and subjected to column chromatography with 10-50 % CH_2Cl_2 /hexane. The alkene




25

was obtained in 67 % yield. ^1H NMR (CDCl_3) δ 1.74-1.97 (m, 12H), 2.60 (br s, 1H), 3.24 (br s, 1H), 3.29 (s, 3H), 7.18-7.26 (m, 2H), 7.38-7.42 (m, 1H), 7.47 (s, 1H).

30 A solution of 2.02 g of the bromoalkene and 1.68 mL of

2. Synthesis of Benzothiazoleboronic acid Derivatives as Signalling Compounds for Fluorescent Detection.



2a) $R^9 = \text{COOCH}_3$

2b) $R^9 = \text{CONH}_2$

2c) $R^9 = \text{CN}$

5

a) Methyl 6-hydroxybenzothiazole-2-carboxylate was prepared by esterification of 6.0 g of the acid with HCl gas in 100 mL of methanol. Product (5.05 g, 79 %) was isolated from the reaction by filtration, air drying and washing with CH_2Cl_2 followed by hexanes. ^1H NMR ($\text{DMSO}-d_6$) δ 3.94 (s, 3H), 7.13 (dd, 1H), 7.50 (d, 1H), 8.01 (d, 1H).

The hydroxy group of the ester was converted to the triflate ester by reacting 5.0 g with 13.0 g of N-phenyltrifluoromethanesulfonimide in 100 mL of CHCl_3 and 12 mL of triethylamine at reflux for 1 h under Ar. The reaction mixture was cooled, diluted to 250 mL with CHCl_3 and extracted sequentially with 20 % aq. citric acid, water, and satd. aq. NaHCO_3 . The organic layer was dried and evaporated. The product triflate (6.2 g, 76%) was isolated by silica gel column chromatography with 25-100 % CH_2Cl_2 /hexanes. ^1H NMR (CDCl_3) δ 4.12 (s, 3H), 7.51 (dd, 1H), 7.95 (d, 1H), 8.32 (d, 1H).


A reaction mixture containing 6.0 g of the triflate, 5.4 g of bis(pinacolato)diboron, 0.40 g of $\text{Pd}(\text{OAc})_2$, 0.93 g of triphenylphosphine and 8.1 g of CsF in 80 mL of anhydrous acetonitrile was refluxed under Ar for 20 min. The mixture was cooled, diluted with acetonitrile and filtered through Celite. The filter bed was washed with CH_2Cl_2 and EtOAc. The combined filtrates were adsorbed onto dry silica and the material subjected to column chromatography with 10-30%

ethyl acetate/hexanes. Product 2a was obtained (3.55 g, 60%) by evaporating pooled fractions containing the product, washing the solid with hexane and air drying. ^1H NMR (CDCl_3) δ 1.38 (s, 12H), 4.09 (s, 3H), 7.98 (d, 1H), 8.22 (d, 1H), 8.47 (s, 1H).

b) Signalling compound 2a (3.05 g) was dissolved in 100 mL of methanol. Anhydrous NH_3 gas was bubbled through the solution causing a rise on temperature. Bubbling was continued for 10 min. The reaction flask was capped and the solution allowed to stand over night. The solvent was then evaporated to produce signalling compound 2b as a white solid 2.91 g, 100 %. ^1H NMR (CDCl_3) δ 1.38 (s, 12H), 5.86 (bs, 1H), 7.34 (bs, 1H), 7.9 (d, 2H), 8.08 (d, 2H), 8.47 (s, 1H).

c) Signalling compound 2b was converted to nitrile 2c with POCl_3 . Compound 2b (0.50 g) was added to 10 mL of POCl_3 . The mixture was stirred under a stream of Ar at 95 $^\circ\text{C}$ for 4 h. The mixture was cooled and evaporated to dryness. The residue was taken up in acetonitrile and poured into ice-cold saturated NaHCO_3 solution. A tan solid formed and was collected by filtration, washed with water and air dried. Compound 2c, 0.40 g, 85 % yield. ^1H NMR (CDCl_3) δ 1.38 (s, 12H), 8.04 (d, 1H), 8.21 (d, 1H), 8.46 (s, 1H).

3. Synthesis of Coumarinboronic ester Derivatives as Signalling Compounds for Fluorescent Detection.




7-Hydroxy-4-methylcoumarin was converted to the triflate ester by reacting 5.0 g with 15.2 g of N-phenyltrifluoromethanesulfonimide in 200 mL of CHCl_3 and 20 mL of triethylamine at reflux for 5 h under Ar. During the reflux period an additional 3.0 g of N-phenyltrifluoromethanesulfonimide was added. The reaction mixture was cooled over night and extracted with 2 x 200 mL of 10% citric acid followed by water wash. The organic layer was dried over Na_2SO_4 and evaporated. The clear colorless oil was crystallized from ether yielding the triflate ester, 8.72 g, 81%. ^1H NMR (CDCl_3) δ 2.47 (s, 3H), 6.37 (s, 1H), 7.23-7.29 (m, 2H), 7.7 (d, 1H).

A reaction mixture containing 2.0 g of the triflate, 2.47 g of bis(pinacolato)diboron, 164 mg of $\text{Pd}(\text{OAc})_2$, 372 mg of triphenylphosphine and 4.6 mL of triethylamine in 30 mL of anhydrous acetonitrile was refluxed under Ar for 4 h. The mixture was cooled and filtered through silica gel with acetonitrile. The filtrate was adsorbed onto dry silica and the material subjected to column chromatography with 0-50% ethyl acetate/hexanes. Since the product was determined to be impure it was subject to a second column using 50-100% CH_2Cl_2 /hexanes. Product 3a was obtained (326 mg, 17%). ^1H NMR (CDCl_3) δ 1.37 (s, 12H), 2.45 (s, 3H), 6.33 (s, 1H),

7.58 (d, 1H), 7.70 (d, 1H), 7.74 (s, 1H).

A reaction mixture containing 251 mg of the triflate, 430.6 mg of bis(neopentyl glycolato)diboron, 23 mg of $\text{Pd}(\text{Oac})_2$, 49 mg of triphenylphosphine, 0.60 mL of triethylamine and 15 mL of anhydrous acetonitrile was refluxed under Ar for 6 h and cooled over night. The mixture was filtered through silica gel with acetonitrile. The solvent was evaporated and the residue subjected to column chromatography with 0-50% CH_2Cl_2 /hexanes. Product 3b was obtained (79 mg, 36%) containing a small amount of neopentyl glycol as an impurity. ^1H NMR (CDCl_3) δ 1.04 (s, 6H), 3.80 (s, 4H), 6.31 (s, 1H), 7.57 (d, 1H), 7.69 (d, 1H), 7.74 (s, 1H).

4. Synthesis of Pyreneboronic acid Derivatives as Signalling Compounds for Fluorescent Detection.



A solution of 1.03 g of 2-bromopyrene in 30 mL of anh. THF was cooled to -78°C under Ar. A solution of 1.50 mL of 2.5 M n-BuLi in hexanes was added dropwise causing a yellow precipitate to form. The reaction mixture maintained at -78°C for another 30 min before dropwise addition of 0.42 mL of trimethylborate. After an additional 2 h at -78°C , the reaction mixture was warmed to room temperature. The reaction was quenched with 5 mL of water. Addition of

CH_2Cl_2 caused formation of a precipitate of 381 mg of nearly pure 4a. The filtrate was evaporated and the residue chromatographed with 20-30 % EtOAc/hexanes to provide an additional 76 mg of 4a. ^1H NMR ($\text{DMSO}-d_6$) δ 7.90

5 (d, 1H), 8.06-8.42 (m, 7H), 9.00 (d, 1H).

A solution of 4.00 g of 2-bromopyrene, 3.97 g of bis(pinacolato)diboron, 4.19 g of KOAc, and 0.29 g of $\text{Pd}(\text{dppf})\text{Cl}_2$, in 30 mL of anh. DMSO was put under Ar and heated at 85 °C for ca. 20 h. The mixture was cooled,

10 poured into 200 mL of water and extracted with 3 x 100 mL of CH_2Cl_2 . The combined CH_2Cl_2 extracts were washed with water, dried over Na_2SO_4 and evaporated. The residue was chromatographed on silica using 5-30 % EtOAc/hexanes.


Boronate ester 4b was obtained 3.61 g (78 %) as a light
15 yellow solid. ^1H NMR (CDCl_3) δ 1.49 (s, 12H), 7.97-8.22 (m, 7H), 8.54 (d, 1H), 9.07 (d, 1H).

Compound 4b was converted to compound 4a by reacting 1.0 g of 4b with 1.96 g of NaIO_4 and 0.52 g of NH_4OAc in 100 mL of 50 % aq. acetone with stirring for 5 days. The solution
20 was filtered, the solids washed with 100 mL of acetone and the filtrates combined and evaporated. Compound 4a was isolated from the residue by silica gel chromatography with 30-100 % EtOAc/hexanes followed by 20 % MeOH/EtOAc in 86 % yield.


25

5. Synthesis of Phenanthreneboronic acid Derivatives as Signalling Compounds for Fluorescent Detection.

5



5a



5b


A solution of 2.00 g of 9-bromophenanthrene, 2.17 g of bis(pinacolato)diboron, 2.29 g of KOAc, and 0.19 g of Pd(dppf) Cl_2 , in 10 mL of anh. DMSO was put under Ar and heated at 85 °C over night. The mixture was cooled, poured into 100 mL of water causing a white precipitate to form. The mixture was extracted with 2 x 100 mL of EtOAc. The combined EtOAc extracts were washed with water (emulsion!), dried over Na_2SO_4 and evaporated. The residue was chromatographed on silica using 10 % EtOAc/hexanes. Boronate ester 5b was obtained 2.27 g (96 %) as a light yellow solid. ^1H NMR (CDCl_3) δ 1.46 (s, 12H), 7.55-7.70 (m, 4H), 7.94 (d, 1H), 8.39 (s, 1H), 8.66-8.73 (m, 2H), 8.81-8.85 (m, 1H).

Compound 5b was converted to compound 5a by reacting 2.07 g with 4.37 g of NaIO_4 and 1.15 g of NH_4OAc in 100 mL of 50 % aq. acetone with stirring for 6 days. The solution was filtered, the solids washed with 50 mL of acetone and the filtrates combined and evaporated. Compound 5a (1.06 g, 70 %) was isolated from the residue by silica gel chromatography with 30-100 % EtOAc/hexanes followed by 20 % MeOH/EtOAc and then washing the solid product in CHCl_3 . ^1H NMR ($\text{DMSO}-d_6$) δ 7.64-7.74 (m, 4H), 8.12 (d, 1H), 8.45-8.92

(m, 2H), 8.70 (s, 1H), 9.36 (d, 1H).

6. Synthesis of Anthraceneboronic acid Derivatives as
Signalling Compounds for Fluorescent Detection.

5




10

6

A solution of 2.00 g of 9-bromoanthracene, 2.17 g of bis(pinacolato)diboron, 2.29 g of KOAc, and 0.19 g of Pd(dppf)Cl₂, in 10 mL of anh. DMSO was put under Ar and heated at 85 °C for 24 h. The mixture was cooled, poured
15 into 100 mL of water causing a white precipitate to form. The mixture was extracted with 3 x 100 mL of EtOAc. The combined EtOAc extracts were washed with water (2 x 100 mL), dried over Na₂SO₄ and evaporated. The residue was chromatographed on silica using 10-25 % EtOAc/hexanes.
20. Boronate ester 6 was obtained 1.29 g (54 %) as a white solid. ¹H NMR (CDCl₃) δ 1.58 (s, 12H), 7.41-7.51 (m, 4H), 7.99 (d, 2H), 8.44 (d, 2H), 8.48 (s, 1H).

7. Synthesis of Bis(benzothiazole)boronic acid Derivatives
as Signalling Compounds for Fluorescent Detection.



A solution of 2.0 of 6-bromo-2-cyanobenzothiazole, and 1.0 mL of 2-aminothiophenol in 100 mL of methanol was stirred under Ar for several days. The white solid was collected by filtration and washed sequentially with methanol, 2-propanol and hexane. Air drying left 2.71 g (93 %)

15

of 6-bromo-2,2'-bis(benzothiazolyl). The bromide (0.50 g), 0.402 g of bis(pinacolato)diboron, 0.424 g of KOAc, and 35.3 mg of Pd(dppf)Cl₂, in 10 mL of anh. DMSO was put under Ar and heated at 85 °C over night.

20

The mixture was cooled, poured into 200 mL of water causing a brown precipitate to form. The solid was collected by filtration and washed sequentially with water, 2-propanol and hexane. Air drying left 0.356 g (63 %) of compound 7b.

¹H NMR (CDCl₃) δ 1.39 (s, 12H), 7.47-7.59 (m, 2H), 7.98 (t, 2H), 8.13-8.18 (m, 2H), 8.47 (s, 1H).

25


The bromide (0.50 g), 0.358 g of bis(neopentylglycolato)diboron, 0.424 g of KOAc, and 35.3 mg of Pd(dppf)Cl₂, in 10 mL of anh. DMSO was put under Ar and heated at 85 °C over night. The mixture was cooled and filtered and the solids washed with acetone. This solid was taken up in 125

30

mL of CH_2Cl_2 and passed through a plug of Celite to remove residual Pd. Evaporation of solvent produced 375 mg of compound 7c (68 %). ^1H NMR (CDCl_3) δ 1.06 (s, 6H), 3.83 (s, 4H), 7.46-7.59 (m, 2H), 7.96-8.01 (m, 2H), 8.12-8.18 (m, 2H), 8.46 (s, 1H).

Attempted dissolution of 7c in D_2O for NMR analysis caused rapid hydrolysis to compound 7a. The filtrate from the initial reaction was poured into 200 mL of water causing a brown precipitate to form. The solid was collected by filtration and washed sequentially with 2-propanol and hexane to provide compound 7a. ^1H NMR ($\text{DMSO}-d_6$) δ 7.57-7.68 (m, 2H), 8.02 (d, 1H), 8.14-8.28 (m, 3H), 8.34 (s, 1H), 8.59 (s, 1H).

8. Synthesis of Benzothiazoleboronic acid Derivative as Signalling Compound for Luminescent Detection.




8

A solution of 390 mg of compound 2c in 25 mL of methanol was sparged with Ar. An Ar-sparged solution of 290 mg of D-cysteine hydrochloride in 5 mL of water adjusted to pH 8 with NaHCO_3 was added by pipet. An additional 5 mL of water was added to the solution. Condensation of the nitrile and cysteine was immediate so the methanol was evaporated and the remaining solution diluted to 100 mL with 20 % conc. HCl. The white precipitate was collected by filtration, washed with water and air dried. Compound 8 was obtained 381 mg, 91 % yield. ^1H NMR (CDCl_3) δ 3.68-3.84 (sextet, 2H),

5.46 (t, 1H), 7.99 (d, 1H), 8.12 (d, 1H), 8.33 (s, 2H),
8.54 (s, 1H).

9. Synthesis of an Indoleboronic Acid Signalling Compound.

5



9


3-Bromoindole is converted to the p-toluenesulfonamide
10 derivative by reaction with p-toluenesulfonyl chloride and
triethylamine. Alternatively, indole can be protected as
the SEM derivative by reaction with (Me₃SiOCH₂CH₂OCH₂Cl)
SEMCl and NaH. The N-protected indole is brominated at the
3-position by reaction with pyridinium tribromide or N-
15 bromosuccinimide. The N-protected 3-bromoindole derivative
is converted to the 3-boronic acid derivative by metal-
halogen exchange with t-BuLi in deoxygenated anh. THF at-
78 °C followed by reaction with trimethyl or triisopropyl
borate and aqueous hydrolysis. Finally the protecting group
20 is removed. Sulfonamides are hydrolyzed by reaction with
KOH in ethanol; SEM groups are removed with LiBF₄ in CH₃CN
followed by aq. NaOH.

10. Chemiluminescent Detection of Hydrogen Peroxide with
25 Boronic Acid-Substituted Dioxetane 1. The dioxetane
signalling compound of Example 1 was prepared in reagent
compositions containing chemiluminescence enhancers and
reacted with peroxide at 37 °C. The solutions contained 1
mM dioxetane and 1 mg/mL of enhancer in 0.3 M tris buffer,
30 pH 9.35. Urea peroxide was added to achieve a final

concentration of 10^{-5} M.

	<u>Enhancer</u>	<u>t max</u>	<u>I_{max} (RLU)</u>	<u>Background</u>	<u>I/B</u>
	None	5	10.75	0.282	38
	CTAB	11.5	49.2	0.81	61
5	Plus [†]	25	1067	12.1	88
	TB [‡]	11.5	1001	5.9	170
	3TB/TO [§]	15	>10000	123	> 100

CTAB - cetyltrimethylammonium bromide



§ - poly(vinylbenzyltributylphosphonium chloride)
copolymer, Compound TB in U.S. Patent 5,393,469

‡ - poly(vinylbenzyltributyl(trioctyl)phosphonium
15 chloride) copolymer, in U.S. 5,393,469

11. Sensitivity of Detection of Hydrogen Peroxide with a Boronic Acid-Substituted Dioxetane.

A reagent composition (90 μ L) containing 1 mM dioxetane signalling compound of
20 Example 1 in 0.3 M tris buffer, pH 9.35 and 1 mg/mL of a
poly(vinylbenzyltributylphosphonium chloride) copolymer,
Compound TB in U.S. Patent 5,393,469, was reacted with
various concentrations of peroxide (10 μ L) over the range
10 mM to 1 μ M. Chemiluminescence intensity was measured
25 after 4.5 min on triplicate samples. Figure 1 shows a plot
of the corrected chemiluminescence intensity as a function
of the peroxide concentration in the reaction. The signal
from the reaction solution with a final peroxide
concentration of 1 mM was ca. 3000 times that of the blank.

12. Fluorescent Detection of Hydrogen Peroxide with 2-Naphthylboronic Acid. A solution of 1 mM 2-naphthylboronic acid was prepared in 0.2 M 221 buffer, pH 9.6. A 0.1 M stock solution of urea peroxide was prepared in deionized water.


5 Ten-fold serial dilutions of the peroxide solution were prepared down to 1×10^{-9} M. Reaction solutions containing 100 μ L of 2-naphthylboronic acid solution and 100 μ L of each peroxide dilution were prepared in 1 cm cuvettes. Fluorescence excitation and emission maxima for the product
10 were 380 nm and 465 nm, respectively.

	<u>Final [Peroxide] M</u>	<u>IFL (10 min)</u>
	5×10^{-4}	103.1
	5×10^{-5}	15.85
	5×10^{-6}	5.76
15	5×10^{-7}	4.74
	Blank	4.60

13. Fluorescent Detection of Glucose Oxidase with 2-Naphthylboronic Acid. Dilutions of glucose oxidase
20 containing from 4.5×10^{-14} to 1.5×10^{-17} moles of enzyme in 3 μ L of water were placed in triplicate white microwells. A solution of glucose (50 μ L of 0.1 M) in 10 mM tris buffer, pH 7.0 was added to each well. The wells were incubated at 25 $^{\circ}$ C for 30 min. Then 50 μ L portions of the reagent of
25 Example 15 were added to each well. Fluorescence was measured after 15 min. Figure 2 depicts the linear response of fluorescence to glucose oxidase over the range of the assay.

14. Fluorescent Detection of Glucose with 2-Naphthylboronic Acid. Dilutions of glucose containing between 5×10^{-8} and 5×10^{-11} moles of glucose in 3 μL of water were placed in triplicate white microwells. Then 100 μL portions of the reagent of Example 12 also containing 3×10^{-7} M glucose oxidase were added to each well. Fluorescence was measured after 30 min. Figure 3 depicts the linear response of fluorescence to glucose oxidase over the range of the assay.

15. Fluorescent Detection of Hydrogen Peroxide with a Bis(benzothiazolylboronate Ester. 50 μL of a 1 mM solution of the bis(benzothiazolyl)boronic acid ester 7b




7b

in 0.2 M 2-amino-2-methyl-1-propanol buffer, pH 9.6 was treated with aqueous solutions containing hydrogen peroxide. Fluorescence excitation and emission spectra shown in Figure 4 were measured before and after addition of 0.1 mM urea peroxide with a Jobin Yvon/SPEX Fluoroskan spectrofluorometer. Fluorescence excitation and emission maxima for the product were 419 nm and 571 nm, respectively.

16. Fluorescent Detection of Hydrogen Peroxide with 4-Methylcoumarin-7-boronic Acid. A 50 μL solution of 0.5 mM 4-methylcoumarin-7-boronic acid pinacol ester in 0.3 M tris buffer, pH 9.3 was prepared. Urea peroxide was added to

achieve a concentration of 3 mM. The grow-in of fluorescence at 440 nm is shown in figure 5.



Fluorescence excitation and emission maxima for the product were 360 nm and 440 nm, respectively.

17. Fluorescent Detection of Urea Peroxide with Other Fluorescent Signalling Compounds. Procedure: A 0.1 mM solution of each of the signalling compounds listed in the table was prepared in 0.3 M tris buffer, pH 9.35. A 3 mL portion of each solution was added to a 1 cm cuvette and the emission spectrum recorded. Urea peroxide was added to each solution to achieve a concentration of 1 mM and the resulting solutions allowed to react for 15 min. The emission spectrum was recorded at this point and again after 24 hours. In each case the fluorescence of the phenolic product reached maximum intensity in 15 min and remained constant at 24 hours.

Compound	λ_{ex} (nm)	λ_{em} (nm)	Intensity (RLU)
4a	335	480	1×10^5
4b	335	480	1×10^5
5a	400	450	2×10^6
5b	400	450	2×10^6

The starting boronic acid or ester signalling compounds were negligibly fluorescent under the conditions of measurement. The maximal emission wavelengths shifted ca.

100 nm on conversion of the signalling compound to the

corresponding phenolic product.

18. Colorimetric Detection of Hydrogen Peroxide with 4-

Nitrophenylboronic Acid. A 20 mM stock solution of 4-

5. nitrophenylboronic acid was prepared in ethanol and diluted 1:100 in 0.3 M tris buffer, pH 9.35 to prepare a 0.2 mM working solution A. A 1.0 M stock solution of urea peroxide was prepared in deionized water. Ten-fold serial dilutions were prepared down to 1×10^{-9} M from the stock solution.
- 10 Reaction solutions containing 3.0 mL of solution A and 10 μ L of each peroxide dilution were prepared in 1 cm cuvettes and the absorbance at 405 nm measured in each cuvette as a function of time.

<u>Absorbance</u>					
15	<u>Final [Peroxide] M</u>	<u>10 min</u>	<u>20 min</u>	<u>60 min</u>	<u>24 hours</u>
	3.3×10^{-3}	0.183	0.500	1.393	3.57
	3.3×10^{-4}	0.028	0.062	0.186	1.839
	3.3×10^{-5}	0.011	0.013	0.028	0.249
	3.3×10^{-6}	0.025	0.024	0.027	0.049
20	3.3×10^{-7}	0.013	0.012	0.013	0.019

19. Bioluminescent Detection of Hydrogen Peroxide with

Boronic Acid-Substituted Compound 8. Reaction of compound 8 with hydrogen peroxide produced D-luciferin. Reaction of


25 the generated D-luciferin with ATP, Mg^{+2} salt and luciferase produced the characteristic greenish bioluminescence.

20. Dual Chemiluminescent/Colorimetric Detection of


- 30 Hydrogen Peroxide. A solution is prepared containing the

indoleboronic acid signalling compound of example 9 (0.1 mM) in 0.1 M diethanolamine buffer, pH 10. A 1.0 M stock solution of hydrogen peroxide is prepared in deionized water. Ten-fold serial dilutions are prepared down to 1×10^{-9} M from the stock solution. Reaction solutions containing 2 mL of indoleboronic acid solution and 100 μ L of each peroxide dilution or water for a blank were prepared in 1 cm cuvettes. A 100 μ L aliquot is transferred to a luminometer tube and chemiluminescence measured after 2 min. Absorbance of the reaction solutions is measured at a suitable wavelength, e.g. 600 nm.

21. Synthesis of Additional Dioxetaneboronic Acids.




Dioxetanes 10a-b are prepared by a process analogous to the one described in example 1. The known compounds 5-chloro-2-adamantanone and either methyl 4-chloro-3-bromobenzoate (for 2a) or methyl 3-bromobenzoate (for 2b) are coupled with the reagent prepared from TiCl_3 and LiAlH_4 . The bromoalkene is converted to the boronic acid alkene by reaction with B(Oi-Pr)_3 and $n\text{-BuLi}$ in THF followed by hydrolysis. The boronic acid alkene is photooxygenated by irradiation with a Na lamp under a stream of O_2 with methylene blue sensitizer to yield dioxetane 2a or 2b.



11a


5 Dioxetane 11a is prepared by a process analogous to the one described in example 1. The known diisopropyl ketone and methyl 3-bromobenzoate are coupled with the reagent prepared from TiCl_3 and LiAlH_4 . The bromoalkene is converted to the boronic acid alkene by reaction with B(Oi-Pr)_3 and $n\text{-BuLi}$ in THF followed by hydrolysis. The boronic acid alkene is photooxygenated by irradiation with a Na lamp under a stream of O_2 with methylene blue sensitizer to yield dioxetane 11.



11b

15 Dioxetane 11b was prepared as follows. A solution of 12.0 g of 3-bromo-4-chlorobenzoic acid and 1 mL of H_2SO_4 in 500 mL of methanol was heated at reflux for 9 h. Another 1 mL of H_2SO_4 was added and reflux continued for several more hours until TLC analysis indicated complete consumption of the starting acid. The solvent was evaporated, the yellow solid taken up in 100 mL of CH_2Cl_2 , and extracted sequentially with water, satd. NaHCO_3 , and satd. NaCl . The organic layer was dried over Na_2SO_4 , and evaporated yielding 12.54 g of the ester as a pale yellow solid. ^1H NMR (CDCl_3) δ 3.928 (s, 3H), 7.5 (d, 1H), 7.9 (d, 1H), 8. (d, 1H).

A mixture of 2.00 g of the ester, 0.1637 g of Pd(dppf)Cl₂, 2.240g of bis(pinacolato)diboron, and 2.36 g of KOAc was put under Ar followed by addition of 20 mL of anh. DMSO. The mixture was further treated by repeated
5 vacuum evaporation and Ar purging. The mixture was heated at 85 °C over night. TLC analysis (10% EtOAc/hex) indicated the starting material to be consumed. The reaction mixture was poured into 125 mL of water and extracted with 3 x 75 mL of ethyl acetate. The combined organic extracts were
10 washed with water, dried and evaporated to dryness leaving a brown oil. The product was further purified by silica gel chromatography eluting with 10-30 % ethyl acetate in hexane. The boronate ester-substituted ester of the formula




was obtained as 1.99 g of white solid in 84 % yield. ¹H NMR (CDCl₃) δ 1.383 (s, 12H), 3.916 (s, 3H), 7.41-7.43 (d, 1H),
20 7.97-8.01 (t, 1H), 8.34 (s, 1H).

A mixture of 8.06 g of TiCl₃ and 0.99 g of LiAlH₄ was formed in 100 mL of cooled anh. THF under Ar by slow addition of the LiAlH₄ to the reaction system. A black mixture resulted. The mixture heated to reflux and
25 triethylamine (7.5 mL) was added. After 1 hour at reflux a solution of 1.55 g of the ester of the previous step and 1.79 g of 2,4-dimethyl-3-pentanone in 20 mL of anh. THF was added dropwise. The reaction mixture was cooled and carefully poured into a solution of 1 L of water and 70 mL
30 of triethylamine. The mixture was extracted with 3 x 400 mL


of ethyl acetate and the aq. Layer discarded. The ethyl acetate extracts were combined, dried over Na_2SO_4 , filtered and evaporated to dryness leaving 1.43 g of a colorless oil. The oil was subjected to column chromatography with 0-
 5 1 % ethyl acetate/hexane producing 0.79 g of a fraction containing the product alkene. ^1H NMR (CDCl_3) δ 0.91 (d, 6H), 1.24 (d, 6H), 1.38 (s, 12H), 2.29-2.40 (m, 2H), 3.17 (s, 3H), 7.19-7.23 (dd, 1H), 7.32 (d, 1H), 7.55 (d, 1H).

10



A solution of the boronate ester-substituted alkene
 15 (50.7 mg) was photooxygenated in 10 mL of CD_3OD containing 50.3 mg of polymer-bound Rose Bengal in an ice bath for 20 min. Oxygen gas was bubbled through the solution while it was irradiated with 1000 W Na lamp through a Kapton optical cutoff filter. The beads were filtered off, the solvent was
 20 evaporated and the residue stored at -20°C . ^1H NMR (CDCl_3) δ 0.91 (d, 6H), 1.24 (d, 6H), 1.38 (s, 12H), 2.29-2.41 (m, 2H), 3.17 (s, 3H), 7.19-7.23 (dd, 1H), 7.32 (d, 1H), 7.55 (d, 1H).


25




12

Dioxetane 12 is prepared by a process analogous to the one
 30 described in example 1. The known compound 3-bromo-4-


chlorobenzoic acid is esterified in 2,2,2-trifluoroethanol with acid catalysis to prepare the trifluoroethyl ester. Adamantanone and 2,2,2-trifluoroethyl 3-bromo-4-chlorobenzoate are coupled with the reagent prepared from TiCl_3 and LiAlH_4 . The bromoalkene is converted to the boronic acid alkene by reaction with $\text{B}(\text{O}i\text{-Pr})_3$ and $n\text{-BuLi}$ in THF followed by hydrolysis. The boronic acid alkene is photooxygenated by irradiation with a Na lamp under a stream of O_2 with methylene blue sensitizer to yield dioxetane 12..



Dioxetane 13 is prepared by a process analogous to that disclosed in EP 0779293A1 depicted below. Briefly, 3-bromobenzoic acid is esterified with 2,2,4,4-tetramethyl-1,3-propanediol to produce the ester (i). The alcohol function is oxidized with pyridinium chlorochromate (PCC) to form (ii). Reductive coupling of the ketone and ester groups with $\text{TiCl}_3/\text{LiAlH}_4$ produces the cyclic vinyl ether (iii). Conversion to the boronic acid is done with triisopropyl borate, followed by hydrolysis as described above. The double bond is photooxygenated as above to produce dioxetane 13.




15



20 Dioxetane 14 is prepared by a process analogous to the one described in example 1. The known ketone tricyclo[7.3.2.0^{2,7}]tridec-2,7-ene-13-one (U.S. Patent 6,461,876) and methyl 4-chloro-3-bromobenzoate are coupled with the reagent prepared from TiCl_3 and LiAlH_4 . The bromoalkene is converted to the boronic acid alkene by reaction with $\text{B}(\text{Oi-Pr})_3$ and $n\text{-BuLi}$ in THF followed by hydrolysis. The boronic acid alkene is photooxygenated by irradiation with a Na lamp under a stream of O_2 with methylene blue sensitizer to yield dioxetane 14.

30




5


Dioxetane 15 is prepared by a process analogous to the one described in example 1. The known compound 3-bromo-4-chlorobenzoic acid is esterified in 2,2,2-trifluoroethanol with acid catalysis to prepare the trifluoroethyl ester.

10 Adamantanone and 2,2,2-trifluoroethyl 3-bromo-4-chlorobenzoate are coupled with the reagent prepared from TiCl_3 and LiAlH_4 . The bromoalkene is converted to the boronic acid alkene by reaction with $\text{B}(\text{O}i\text{-Pr})_3$ and $n\text{-BuLi}$ in THF followed by hydrolysis. The boronic acid alkene is
15 photooxygenated by irradiation with a Na lamp under a stream of O_2 with methylene blue sensitizer to yield dioxetane 15.


22. Synthesis of Benzothiazoleboronic Acid Derivatives as
20 Signalling Compounds for Luminescent Detection.



15a



15b



15c


25
30 A solution of 1.0 g of compound 2c and 620 mg of DL-

penicillamine was prepared in 30 mL. THF. To this was added 2.0 mL of triethylamine was added followed by 3.0 mL of water. The solution was stirred for 1 h while being sparged with Ar. The solvent was evaporated, the residue taken up in CH_2Cl_2 and the solution dried on MgSO_4 . Evaporation of the solvent yielded the triethylamine salt of 15a. The free acid was liberated by taking the residue up in CH_2Cl_2 , and washing the solution with 5% aq. citric acid followed by water, drying on MgSO_4 and evaporation of solvent. Compound 15a was obtained 1.34 mg, 92 % yield. ^1H NMR (CDCl_3) δ 1.379 (s, 12H), 1.581 (s, 3H), 1.910 (s, 3H), 4.970 (s, 1H), 7.957 (d, 1H), 8.140 (d, 1H), 8.432 (s, 1H).

Treatment of the boronate ester with aq. HCl as in example 8 produces the boronic acid derivative 15b.

Compound 15a (1.0 g) and 0.44 g of carbonyl diimidazole (CDI) were added to 30 mL of anh. CH_3CN . The reaction was stirred under Ar for 1 h before adding 0.5 mL of thiophenol. After stirring over night the yellow precipitate was filtered, washed with CH_3CN and air dried producing 0.86 g of 15c (70%). ^1H NMR (CDCl_3) δ 1.383 (s, 12H), 1.577 (s, 3H), 1.845 (s, 3H), 4.939 (s, 1H), 7.451 (s, 5H), 7.962 (d, 1H), 8.152 (d, 1H), 8.440 (s, 1H).

Attempted preparation of the phenyl ester of 15a by esterification with phenol and CDI produced instead



the oxidation product. This compound is useful as a fluorescent signalling compound. ^1H NMR (CDCl_3) δ 1.383 (s,

12H), 2.482 (s, 3H), 2.539 (s, 3H), 7.954 (d, 1H), 8.116 (d, 1H), 8.420 (s, 1H)

23. Sensitivity of Detection of Hydrogen Peroxide with a

5 Boronate Ester-Substituted Dioxetane. Reagent compositions (90 μ L) containing 0.1 mM dioxetane signalling compound 11b of Example 21 in 0.3 M tris buffer, pH 9.35 and either no enhancer or 1 mg/mL of Plus enhancer, TB or 3TB/TO as in example 10, were reacted with 10 μ L solutions of urea
10 peroxide containing from 1 mM to 0.1 μ M peroxide. Light intensity was measured at the chemiluminescence maximum on triplicate samples. Similarly, a sensitivity study was conducted with the same dioxetane and peroxide concentrations in a reaction buffer 0.2 M 221 buffer, pH
15 9.6 containing 0.88 mM MgCl_2 . In these cases, a linear calibration curve was obtained over the full range tested.

24. Generation of Chemiluminescence with Additional Boron

20 Substituted Dioxetanes. Each of dioxetanes 10a, 10b, 11a, 12, 13, 14 and 15 (10^{-6} moles) are added to separate 100 μ L solutions of hydrogen peroxide (0.1 mM) in water. Sodium hydroxide (10 μ L of 0.1 M) solution is injected causing immediate generation of chemiluminescence from each
25 solution. Alternately the hydrogen peroxide and sodium hydroxide are mixed first and the dioxetane solution injected into the alkaline peroxide solution to generate the chemiluminescence. In another alternative, the peroxide solution is prepared in a buffer solution having a pH of at
30 least 9 and the dioxetane solution injected into the

alkaline buffered peroxide solution to generate the chemiluminescence.

5 The foregoing description and examples are illustrative only and not to be considered as restrictive. It is recognized that modifications of the specific compounds and methods not specifically disclosed can be made without departing from the spirit and scope of the present
10 invention. The scope of the invention is limited only by the appended claims.